

Contents

Zusammenfassung	I
Abstract	V
Abbreviations	XIII
1. Introduction	1
1.1. The cellular response to DNA damage	2
1.1.1. Detection of DNA double strand breaks	2
1.1.2. Transducing the damage signal	3
1.1.3. Cellular response	4
1.1.4. The DNA damage response in pathology and cancer therapy	5
1.2. NF- κ B signaling in presence of DNA double strand breaks	6
1.2.1. DNA double strand break-induced NF- κ B activation	7
1.2.2. NF- κ B-induced cell fates in response to DNA damage	9
1.3. The tumor suppressor p53 and its role in the DNA damage response	11
1.3.1. DNA double strand break-induced p53 signaling	12
1.3.2. Mathematical modeling-driven insights into the regulation of p53 signaling	13
1.3.3. P53-induced cell fates in response to DNA damage and its role in the decision making process	14
1.4. Interplay of the NF- κ B and p53 pathway in context of DNA damage	15
 I. Development and analysis of a dynamical model describing genotoxic NF-κB signaling	 17
2. Motivation	19
3. Results	21
3.1. Modular approach to develop a dynamical model of DNA damage-induced NF- κ B activation	21
3.2. Recognition of DSBs by PARP-1 and the importance of its self-enhanced recruitment	21
3.2.1. Positive feedback of PARP-1 recruitment is not sufficient to reproduce the experimental data	22
3.2.2. Incorporating NAD ⁺ consumption into the model allows to reproduce the data but predicts hyperactivation of PARP-1	24
3.2.3. Model describing recruitment of PARP-1 to periphery of DNA lesions reproduces the data and predicts low numbers of induced DSBs	27
3.3. Recruitment and activation of kinase ATM	30
3.4. Formation of a nuclear complex integrates the signals of the sensor proteins	33
3.5. ATM and IKK γ transfer the DNA damage signal from the nucleus to the cytoplasm	36
3.6. Merging the individual models	38
	IX

3.7. Sensitivity analysis reveals a strong impact of PARP-1 modification on IKK activation	38
3.8. Time-resolved regulation of IKK activity	43
3.8.1. The effect of the irradiation strength on the regulation of IKK activity	43
3.8.2. Inhibition of ATM activity modulates the level of PARylated PARP-1 contributing to signalosome formation	49
3.8.3. Irradiation strength-dependent increase in parPARP levels counteract the effect of ATM inhibition on signalosome formation	52
3.9. Time-resolved impact of parallel signaling on IKK activation	54
4. Discussion	57
II. Deciphering the multifaceted interplay of IKK2 and the p53 pathway using single cell dynamics	63
5. Motivation	65
6. Results	67
6.1. The response of p53 is modulated by the activation status of the NF- κ B pathway	67
6.1.1. Quantified changes in the features of p53 dynamics	68
6.1.2. Consistency of IKK2i-mediated changes in p53 dynamics	69
6.1.3. Specificity of IKK2i-mediated changes in p53 dynamics	71
6.2. Reproducing the heterogeneous dynamics of p53 using subpopulation-based modeling	72
6.2.1. Clustering single cell trajectories reveals ten subpopulations with distinct p53 dynamics	75
6.2.2. Subpopulation-based modeling allows to reproduce the heterogeneous dynamics of p53 upon DSBs	78
6.2.3. Identifying subpopulation-specific parameters driving heterogeneity in the p53 response	80
6.3. IKK2 is predicted to act on multiple processes of the p53 network	82
6.3.1. Peak-wise sensitivity analysis	84
6.3.2. Comparing the results of the sensitivity analysis with the experimental observations	84
6.3.3. Sensitivity analysis of parameter pairs reveals potential interfaces	85
6.3.4. Differential impact of parameters on the first and the consecutive p53 peaks	87
6.4. Parameter inference predicts interactions of IKK2 and p53 network components	88
6.5. Validation of predicted parameter combinations unravels two signaling mechanisms	92
7. Discussion	99
8. Conclusion and future perspectives	107
9. Methods	111
9.1. Quantification of western blot and electro mobility shift assay data	111
9.2. Parameter inference	112
9.3. Profile Likelihood Estimation	114
9.4. Akaike weights	114

9.5. Adapting the Data2Dynamics toolbox	115
9.6. Sensitivity analysis of overall model	116
9.7. Determining the area under the curve	116
9.8. Processing single cell time lapse data and quantifying features of dynamics	117
9.8.1. Smoothing of single cell trajectories	117
9.8.2. Filter for trajectories reaching detection limits	117
9.8.3. Quantifying features of single cell dynamics	117
9.9. Clustering of single cell trajectories	118
9.9.1. Filter for outliers regarding peak number and peak timing	120
9.9.2. Determining dissimilarities between single cell trajectories	120
9.9.3. Clustering	122
9.9.4. Evaluation of different cluster settings	128
9.10. Subpopulation-based modeling	129
9.10.1. Defining and implementing subpopulation-specific parameters	129
9.10.2. Fit of model pool to subpopulations	131
9.10.3. Using L1 regularization to identify parameters driving heterogeneity	132
9.11. Sensitivity analysis	137
9.12. Reproducing the perturbation data	138
9.12.1. Processing of perturbation data	138
9.12.2. Allocating trajectories of perturbation data to clusters of calibration data	138
9.12.3. Parameter inference based on perturbation data	143
9.13. Weighted χ^2 value	143
A. Development and analysis of a dynamical model describing genotoxic NF-κB signaling	145
A.1. Overview of experimental data sets	145
A.2. Modeling the auto-modification of PARP-1	146
A.2.1. ODE systems of model variants	146
A.2.2. Parameters of model PM_4	149
A.2.3. Comparison of PARP-1 model variants	151
A.2.4. Additional data for PARP-1 models	152
A.3. Modeling the activation of ATM	155
A.3.1. Model variants describing activation of ATM	155
A.3.2. ODE systems of model variants	155
A.3.3. Parameters of model variant AM_6	160
A.3.4. Comparison of model variants	160
A.3.5. Quantified western blot data of phosphorylated ATM	161
A.3.6. Profile likelihood estimation of model variant AM_5	162
A.4. Modeling the nuclear signalosome formation	162
A.4.1. ODE system of the signalosome formation model	162
A.5. Modeling the TRAF6 complex formation in the cytoplasm	164
A.5.1. Schemes of additional model variants	164
A.5.2. Quantified western blot data calibrate TRAF6 complex formation model variants	166
A.5.3. ODE systems of model variants	167
A.5.4. Parameters of model variant TM_2	169
A.5.5. Overview of model variants	169
A.5.6. PLE for model variant TM_2	170
A.5.7. Simulation of best fit of model variant TM_2	171

A.6. Overall model describing activation of IKK upon DSBs	172
A.6.1. ODE system of the overall model	172
A.6.2. Simulation and experimental data of overall model	174
A.6.3. Parameters of overall model	175
A.6.4. Validation experiments for ATM and PARP-1 inhibition	177
A.6.5. Time-resolved impact of parPARP, pATM and IKK γ on flux of sig- nalosome formation	179
A.6.6. Time-resolved impact of spIKK γ and ATT on flux of IKK activation	182
B. Deciphering the multifaceted interplay of IKK2 and p53 pathway using single cell dynamics	185
B.1. Description and labeling of single cell data sets	185
B.2. Mathematical model of p53 signaling	186
B.2.1. Labels and description of p53 model parameters	186
B.2.2. ODE system of p53 model pool	189
B.3. Additional data of features of dynamics	192
B.3.1. Additional features of dynamics of the reference data set	192
B.3.2. Features of dynamics of the parameter inference data set	193
B.3.3. Features of dynamics of the replicate data set	194
B.3.4. Additional features of dynamics of the inhibitor data set	198
B.4. Subpopulation-based modeling	201
B.5. Sensitivity analyses	204
B.6. Parameter inference based on perturbation data	207
B.7. Validation of predicted parameter combinations	208
List of Figures	209
List of Tables	213
Bibliography	215
Acknowledgements	233